

Masodine[®] 1:3

2.15% w/v concentrate for teat dip & teat spray solution



Economical concentrated formulation, helps to reduce bactericidal contamination. Effective aid in the control of mastitis in dairy herds.

UK Authorised Veterinary Medicine.

Formulated to help reduce bacterial contamination that causes mastitis in dairy herds.

- Reduces the risk of infection by leaving an antiseptic film of Iodine adhering to teats which have been treated by dipping or spraying.
- Use as a post-milking dip or spray at full dilution rate of 1:3.
- May also be used as a pre-milking dip or spray diluted at a rate of 1:7.
- Contains 20% Glycerol & Sorbitol (5% when diluted) to help keep teats soft & supple, & to aid in the prevention & healing of sores & cracks.

TEAT DIP OR SPRAY: (5,000 ppm minimum available Iodine) Dilute 1:3 parts clean water & mix well.

DIPPING: Fill teat cup about two thirds full with the prepared solution. Dip teats of every cow immediately after each has been milked, making sure that the full length of the teat is immersed. Top up with fresh solution as required. Empty & wash out cups after milking.

SPRAYING: Immediately after milking, spray the entire surface of each teat of every cow with prepared solution.

UDDER WASH & CLUSTER DIP: (40 ppm minimum available Iodine)

Dilute 1: 500 parts clean water (20 ml to 10 litres or 1 fl oz in 3 gal).

To help prevent cross infection, individual cloths or disposable paper towels are strongly recommended. Dry teats before milking. Immerse clusters & agitate for at least half a minute. Rinse in clean water before use. Wash & dry udders and teats before milking.

ALTERNATIVE USAGE - PRE-MILKING DISINFECTION: (2,500 ppm minimum available Iodine)

Dilute 1:7 parts clean water & mix well. Ensure udder & teats are clean & dry before pre-dipping.

DIPPING: Fill teat dipping cup 2/3 full with the 1:7 solution. Dip teats of every cow making sure that the full length of the teat is immersed. Allow 15 seconds contact time, then using a single service paper towel, wipe & dry teat thoroughly, before continuing with normal milking routine. Top up with fresh solution as required. Empty & wash out cups after milking.

SPRAYING: Spray the entire surface of each teat of every cow with 1:7 solution. Allow 15 seconds contact time, then using a single service paper towel, wipe and dry teat thoroughly, before continuing with normal milking routine.

5l Order Code: R071EEV2

Pack: 2 x 5 litre (11.57 kg)
Pallet: 60 Pack

25l Order Code: R071KEV

Pack: 25 litre (28.54 kg)
Pallet: 32 Each

200l Order Code: R071NEV

Pack: 200 litre (228.62 kg)
Pallet: 4 Each

1000l Order Code: R071SEV

Pack: 1000 litre (1164 kg)
Each

PRODUCT INFORMATION:

Appearance: Dark brown liquid
Odour: Faint Iodine
pH - undiluted: 4.0
Shelf life: 2 years

COMPOSITION: Contains a blend of iodine, glycerol, sorbitol & nonionic surfactant.

BIODEGRADABILITY: All surfactants used in Evans Vanodine products comply with the current European Regulations concerning biodegradability & protection of the environment.

QUALITY ASSURANCE: This product is manufactured in the U.K. by EVANS VANODINE INTERNATIONAL PLC under an ISO 9001 Quality Management System Cert. No. FM 09535 and an ISO 14001 Environmental Management System Cert. No. EMS 506072 registered by the British Standards Institution.

REVISION DATE:
03/04/18



Evans Vanodine International plc

Brierley Road, Walton Summit, Preston, PR5 8AH, UK
www.evansvanodine.co.uk | Tel: +44 (0)1772 322200 | Fax: +44 (0)1772 626000

